Reflection / Huritau

Emergency Department pandemic preparedness: Putting research into action

Jessica Lockett^{1,2}, MNS, RN, Quality Improvement Advisor

¹Health Quality & Safety Commission, Wellington, Aotearoa New Zealand

Corresponding Author: jcmlockett@gmail.com

Note: Article written when author a Clinical Nurse Educator in the Emergency Department at Capital & Coast DHB, Wellington.

Citation

Lockett, J. (2021). Emergency Department pandemic preparedness: Putting research into action. *Nursing Praxis in Aotearoa New Zealand*, 37(3), 20-21.
<https://doi.org/10.36951/27034542.2021.028>

Abstract

Emergency nurses are experts at responding rapidly to change. Their everyday practice involves constant assessment and prioritisation of a dynamic workload to meet the needs of patients across the spectrum of illness and injury. When COVID-19 arrived in Aotearoa New Zealand in February 2020, staff in hospitals around the country rushed to prepare for the wave of infected patients seen overseas. Emergency nurses' ability to adapt to unanticipated situations whilst working in an overcrowded and resource-constrained system made them important contributors to COVID-19 pandemic planning. Finding myself in the unique position of having just completed novel research into the perspectives of emergency nurses around pandemic preparedness, I felt a responsibility to ensure these perspectives were included in the pandemic response at the busy tertiary emergency department where I worked. Key findings from my prior research included the importance of managing the fear nurses felt about getting sick when caring for patients and of spreading disease to their family, friends, or to other vulnerable patients within their care.

Keywords: COVID-19; emergency department; nursing; pandemic planning

All people living in Aotearoa New Zealand have access to public emergency medical care, but it became apparent early in the pandemic that this right needed to be carefully balanced with protecting emergency department (ED) staff and other patients from the spread of the COVID-19 virus. This reflection focuses on one of the most significant practice changes for nurses; the introduction of COVID-19 screening protocols for all patients and visitors accessing the ED. The early identification and isolation of potentially infectious patients was an important component of the new screening policy and a key strategy that nurses wanted included in future pandemic plans (Lockett et al., 2021).

The burden of screening fell to nurses as the first point of contact for patients on arrival in ED through the triage process. The screening measures underwent many iterations as the COVID-19 situation rapidly evolved and advice from the World Health Organization and the Ministry of Health

frequently changed (Ministry of Health, 2021; World Health Organization, 2021). Triage nurses were initially required to ask about specific international travel, but as COVID-19 spread rapidly internationally, this became unsustainable for nurses tasked with both COVID-19 risk assessment and a general rapid triage assessment. Nurses were concerned they may overlook crucial assessment cues regarding the patient's actual presenting complaint, and that infectious patients may have already transmitted the virus in the waiting room by the time the risk was identified at the point of triage. The ED senior leadership team, of which I was a part of, listened to these concerns and adapted the process to occur before the triage point. This change allowed infectious patients to be diverted from the general waiting room to another entrance where infection control precautions could be maintained throughout. Additionally, screening was extended to visitors, and their details were recorded to allow for retrospective

contact tracing in the event of a positive case. This system preceded by some weeks the introduction of the NZ COVID Tracer app, utilised by the Aotearoa New Zealand government to record the movements of the population (Ministry of Health, 2020).

The separation of infectious and non-infectious patients was a key strategy nurses in my research considered necessary to minimise the risk of disease transmission within the ED. This was based on past instances where vulnerable patients were placed in close proximity to others with infectious diseases, such as influenza, due to environmental constraints of the department or the lack of adequate isolation rooms (Lockett et al., 2021). Although the purpose of this pre-triage screening is to identify and isolate potentially infectious patients, it served a dual purpose that helped to address another fear nurses expressed in my research. Emergency nurses highlighted how challenging the triage process can be and their anxiety around making the right decision. In a pandemic situation, these nurses feared being blamed or censured for not identifying an infectious patient at triage and the impact this oversight could have on staff and other patients. In addressing these fears, the screening tool evolved from an initial three questions into a formalised algorithm that stratified risk based on Ministry of Health case definition advice. This meant the process was objective, rather than a mix of objective and subjective assessment as in routine triage decision-making (Australasian College of Emergency Medicine, 2016). Separating the screening process from the already-challenging triage process also meant that the responsibility for identifying potentially infectious patients was not put solely upon a single individual triage nurse (Lockett et al., 2021).

The screening process, and many other changes introduced during the pandemic, has had a positive impact both on ED daily operations as well as the wider hospital. The screening of patients and visitors at the ED entrance subsequently helps to protect the health and safety of staff in other acute care areas, and reduces the likelihood of an infected patient or visitor accessing a ward environment and spreading infection (Wang et al., 2020). Additionally, as ED was the first area to introduce a formalised screening process, the District Health Board was able to rapidly adapt the ED screening algorithm to be used at visitor entrances when COVID-19 community transmission was at its highest and Aotearoa New Zealand's alert

levels were raised. The underlying motivation behind the introduction of the screening system was the understanding that we could not expect nurses to assuage the fears of the general public during a worldwide pandemic if they themselves did not feel safe in their own workplace. Empowering ED nurses to be involved in the pandemic response was key to ensuring this group of healthcare workers felt heard.

References

- Australasian College of Emergency Medicine. (2016). *Guidelines on the implementation of the Australasian Triage Scale in emergency departments*.
<https://acem.org.au/Content-Sources/Advancing-Emergency-Medicine/Better-Outcomes-for-Patients/Triage>
- College of Emergency Nurses New Zealand. (2016). *College of emergency nurses knowledge & skills framework*.
https://www.nzno.org.nz/groups/colleges_sections/colleges/college_of_emergency_nurses/resources/knowledge_skills_framework
- Lockett, J.C.M., Nelson, K., & Hales, C. (2021). Pre COVID-19 emergency department nurses' perspectives of the preparedness to safely manage influenza pandemics: A descriptive exploratory qualitative study. *Australasian Emergency Care*, 24(4), 280-286.
<https://doi.org/10.1016/j.auec.2021.03.001>
- Ministry of Health. (2021). *COVID-19: Advice for all health professionals*. Retrieved from
<https://www.health.govt.nz/our-work/diseases-and-conditions/covid-19-novel-coronavirus/covid-19-information-health-professionals/covid-19-advice-all-health-professionals>
- Ministry of Health. (2020). *NZ COVID Tracer app released to support contact tracing*. Retrieved from
<https://www.health.govt.nz/news-media/media-releases/nz-covid-tracer-app-released-support-contact-tracing>
- Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., Wang, B., Xiang, H., Cheng, Z., Xiong, Y., Zhao, Y., Li, Y., Wang, X., & Peng, Z. (2020). Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *Journal of the American Medical Association*, 323(11), 1061-1069.
<https://doi.org/10.1001/jama.2020.1585>
- World Health Organization. (2021). *Coronavirus disease (COVID-19) weekly epidemiological update and weekly operational update*. Retrieved from
<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>

Funding: None

Conflicts of interest: None